

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/US2004/034534

International filing date: 20 October 2004 (20.10.2004)

Document type: Certified copy of priority document

Document details: Country/Office: US
Number: 60/512,651
Filing date: 20 October 2003 (20.10.2003)

Date of receipt at the International Bureau: 22 May 2008 (22.05.2008)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

74407

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

May 21, 2008

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE.

APPLICATION NUMBER: 60/512,651

FILING DATE: *October 20, 2003*

RELATED PCT APPLICATION NUMBER: *PCT/US04/34534*

THE COUNTRY CODE AND NUMBER OF YOUR PRIORITY APPLICATION, TO BE USED FOR FILING ABROAD UNDER THE PARIS CONVENTION, IS *US60/512,651*



Certified by

Under Secretary of Commerce
for Intellectual Property
and Director of the United States
Patent and Trademark Office

17638 U.S. PTO
102003

PTO/SB/16 (08-03)

Approved for use through 07/31/2006. OMB 0651-0032

U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

PROVISIONAL APPLICATION FOR PATENT COVER SHEET

This is a request for filing a PROVISIONAL APPLICATION FOR PATENT under 37 CFR 1.53(c).

Express Mail Label No. EL991582016US

22582 U.S. PTO
60/512651
10/20/03

10/20/03

INVENTOR(S)					
Given Name (first and middle (if any))		Family Name or Surname		Residence (City and either State or Foreign Country)	
David N.		Watkins		Baltimore, MD	
Additional inventors are being named on the <u>ONE</u> separately numbered sheets attached hereto					
TITLE OF THE INVENTION (500 characters max)					
Use of Hedgehog Pathway Inhibitors in Small Cell Lung Cancer					
Direct all correspondence to: CORRESPONDENCE ADDRESS					
<input type="checkbox"/> Customer Number: <div style="border: 1px solid black; width: 200px; height: 20px;"></div>					
OR					
<input checked="" type="checkbox"/> Firm or Individual Name		Johns Hopkins University			
Address		100 N. Charles Street			
Address		5th Floor			
City		Baltimore		State	MD
Country		USA		Zip	21201
		Telephone	410-516-8300	Fax	410-516-5113
ENCLOSED APPLICATION PARTS (check all that apply)					
<input checked="" type="checkbox"/> Specification Number of Pages <u>17</u>		<input type="checkbox"/> CD(s), Number _____			
<input type="checkbox"/> Drawing(s) Number of Sheets _____		<input type="checkbox"/> Other (specify) _____			
<input type="checkbox"/> Application Date Sheet. See 37 CFR 1.76					
METHOD OF PAYMENT OF FILING FEES FOR THIS PROVISIONAL APPLICATION FOR PATENT					
<input checked="" type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27. <input type="checkbox"/> A check or money order is enclosed to cover the filing fees. <input type="checkbox"/> The Director is hereby authorized to charge filing fees or credit any overpayment to Deposit Account Number: _____ <input checked="" type="checkbox"/> Payment by credit card. Form PTO-2038 is attached.				FILING FEE Amount (\$) <div style="border: 1px solid black; padding: 5px; text-align: center;">\$80.00</div>	
The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government. <input type="checkbox"/> No. <input checked="" type="checkbox"/> Yes, the name of the U.S. Government agency and the Government contract number are: <u>CA58184</u>					

[Page 1 of 2]

Respectfully submitted,

SIGNATURE Heather Bakalyar
 TYPED or PRINTED NAME Heather Bakalyar, Ph.D.
 TELEPHONE 410-516-8300

Date 10/20/03
 REGISTRATION NO. 45,282
 (if appropriate)
 Docket Number: 4205

USE ONLY FOR FILING A PROVISIONAL APPLICATION FOR PATENT

This collection of information is required by 37 CFR 1.51. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 8 hours to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Mail Stop Provisional Application, Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

If you need assistance in completing the form, call 1-800-PTO-9199 and select option 2.

PROVISIONAL APPLICATION COVER SHEET
Additional Page

PTO/SB/16 (08-03)

Approved for use through 07/31/2006. OMB 0651-0032

U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

Docket Number 4205

INVENTOR(S)/APPLICANT(S)		
Given Name (first and middle [if any])	Family or Surname	Residence (City and either State or Foreign Country)
David M.	Berman	Baltimore, MD
Stephen B.	Baylin	Baltimore, MD
Philip A.	Beachy	Towson, MD

[Page 2 of 2]

Number 2 of 2

WARNING: Information on this form may become public. Credit card information should not be included on this form. Provide credit card information and authorization on PTO-2038.

CERTIFICATE OF EXPRESS MAILING

EXPRESS MAILING LABEL NO.

EL991582016US

I hereby certify that this correspondence (along with any papers referred to as being attached or enclosed) is being deposited with the United States Postal Service as Express Mail, Post Office to Addressee with sufficient postage in a **Flat Rate** envelope addressed to MS Provisional Patent Application, Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450 on the date indicated below:

10/20/03
DATE of Signature
And of Mail Deposit

Cheryl Dixon
Signature

U.S. Provisional Patent Application

JHU Ref. No.: JHU-4205

**Use of Hedgehog Pathway Inhibitors in Small Cell
Lung Cancer**

**Inventors: David N. Watkins, David M. Berman, Stephen B. Baylin and
Philip A. Beachy**

4205

Johns Hopkins University
Licensing and Technology Development

Report of Invention Disclosure Form

MAR 17 2003

This form is to be completed and submitted to the JHU office of Licensing and Technology Development (LTD) by anyone who believes they have developed a new invention. The purpose of this form is to enable LTD to evaluate whether legal protection to the invention will be sought and/or commercialization pursued. In order for this Report of Invention to be processed by LTD, it must be signed and dated by all inventors, and by the JHU Department Director(s) for all departments involved with the development of this invention. LTD can not process this report until it is complete with all necessary signatures found in Sections A, B and/or C. Visit the LTD web site at <http://www.jhu.edu/technology/RptInv.html> for HTML and Word 97 downloadable formats of this form.

INVENTION INFORMATION

Title of Invention:

Use of hedgehog pathway inhibitors in small cell lung cancer

School(s) and Department(s) in which invention was developed: Oncology, Pathology, Molecular Biology&Genetics

Additional inventors: X Yes ☐ No If yes, please complete Additional Inventors section for each inventor.

Lead Inventor Information: [the lead inventor is the primary contact person for LTD]

Name of Lead Inventor:	<u>Watkins</u>	<u>David</u>	<u>Neil</u>	<u>MBBS PhD</u>
	Last	First	Middle	Degree

Title or Position: Research Associate School: Medicine Department: Oncology

Business phone: (410) 955-8506 Business fax: (410) 614-9883 E-mail: nwatkins@jhmi.edu

Business address: Sidney Kimmel Comprehensive Cancer Center At Johns Hopkins
1650 Orleans St, Room 572, Baltimore, MD, 21218.

Interdepartmental address:

Home phone number: 410 662 5717

Home fax number:

Home address: 3120 St Paul St, Apt 414G, Baltimore, MD, 21218

Citizenship: Australian

Social Security Numbe

Are you a Howard Hughes Medical Institute employee or investigator?

☐ Yes ☒ No

Are you a Kennedy Krieger Institute employee or investigator?

☐ Yes ☒ No

LTD Internal Use Only:

REF.

4205

TLA

SHV

Field of Use

ADDITIONAL INVENTOR(S)

Please copy this page for additional inventors as necessary

Name of Inventor: Berman David Monty MD-PhD
Last First Middle Degree

Title or Position: Assistant Professor School: Medicine Department: Pathology

Business phone: (410) 955 -3652 Business fax: (410) 955-9124 E-mail: dberman@jhmi.edu

Business address:
725 N. Wolfe St., PCTB 702
Baltimore, MD 21205
Interdepartmental address:
PCTB 701

Home phone number: 410-366-3276 Home fax number: 419-793-6554

Home address: 3728 Chestnut Ave., Baltimore, MD 21211

Citizenship: USA Social Security Number: _____

Are you an Howard Hughes Medical Institute employee or investigator?

No

Are you a Kennedy Krieger Institute employee or investigator?

No

ADDITIONAL INVENTOR(S)

Please copy this page for additional inventors as necessary

Name of Inventor: Baylin Stephen B. MD
Last First Middle Degree

Title or Position: Prof. School: Johns Hopkins University School of Medicine Department: Oncology

Business phone: 410 955-8506 Business fax: 410 614-9884 E-mail: sbaylin@jhmi.edu

Business address: The Sidney Kimmel Comprehensive Cancer Center at Johns Hopkins
Bunting Blaustein Cancer Research Building
1650 Orleans Street, Suite 541
Baltimore, MD 21231-1000

Home phone number: 410 332-0733 Home fax number:

Home address: 1241 Riverside Avenue
Baltimore, MD 21230

Citizenship: USA Social Security Number:

Are you an Howard Hughes Medical Institute employee or investigator? ☐ Yes ☒ No
Are you a Kennedy Krieger Institute employee or investigator? ☐ Yes ☒ No

ADDITIONAL INVENTOR(S)

Please copy this page for additional inventors as necessary

Name of Inventor: _____ Beachy, Philip A., Ph.D.
Last First Middle Degree

Title or Position: Professor **School:** School of Medicine **Department:** Molecular Biology and Genetics

Business phone: (410)955-1862 **Business fax:** (410) 955-9124 **E-mail:** pbeachy@jhmi.edu

Business address: Philip Beachy, Department of Molecular Biology and Genetics PCTB 714, 725 N. Wolfe Street, Johns Hopkins University School of Medicine, Baltimore MD 21205

Interdepartmental address: Department of Molecular Biology and Genetics, PCTB 714

Home phone number: 410 494-1557 **Home fax number:** _____

Home address: 7818 Chelsea Street, Towson MD 21204

Citizenship: USA **Social Security Number** _____

Are you an Howard Hughes Medical Institute employee or investigator? ☒ Yes ☐ No
Are you a Kennedy Krieger Institute employee or investigator? ☐ Yes ☐ No

INVENTION DESCRIPTION

Describe the invention completely, using the outline given below. Please provide an **Electronic Copy** of the invention disclosure document, references, and abstracts, in Windows format, on CD-Rom or Floppy Disk.

1. **Abstract of the Invention** [In order to assist Licensing and Technology Development with the assessment of this technology, please provide a summary of the invention that should be written to be understood by a wide audience including non-technical individuals].

We have shown that small cell lung cancer (SCLC), a highly aggressive and frequently lethal tumor, uses an embryonic signaling pathway to promote its growth. This signaling pathway is known as the Hedgehog pathway, and its normal function is to regulate organ formation and regulate progenitor cells in embryos and in some adult tissues. SCLC is characterized by abnormal activation of this pathway which can be specifically inhibited by cyclopamine, a naturally occurring compound. Use of cyclopamine specifically inhibits the growth of SCLC, and suggests a potential way to exploit this pathway for novel therapeutic strategies. These strategies could ultimately involve cyclopamine, and other novel inhibitors of Hedgehog pathway, to treat SCLC.

2. **Problem Solved** [Describe the problem solved by this invention]

SCLC is a highly aggressive, frequently lethal form of lung cancer. After initial responses to chemotherapy, the vast majority of patients relapse and die within twelve months. We have discovered that SCLC is vulnerable to specific inhibition of the Hedgehog signaling pathway. Inhibitors of this pathway such as cyclopamine do not act as conventional chemotherapeutic agents and do not produce the same toxicities characteristic of most anticancer drugs. We propose that cyclopamine, and novel inhibitors of Hedgehog signaling, represent potential new therapies that may effectively treat SCLC.

2. **Novelty** [Identify those elements of the invention that are new when compared to the current state of the art]

The novelty of this invention can be summarized as (1) The identification of the Hedgehog signaling pathway in SCLC, (2) The discovery that SCLC growth is dependent on the activity of this pathway, and (3) Specific inhibitors of Hedgehog signaling inhibit the growth of SCLC and represent a potential novel therapy.

4. Detailed Description of the invention:

On a separate page(s), attach a detailed description of how to make and use the invention. The description must contain sufficient detail so that one skilled in the same discipline could reproduce the invention. Include the following as necessary:

- 1- data pertaining to the invention;
- 2- drawings or photographs illustrating the invention;
- 3- structural formulae if a chemical;
- 4- procedural steps if a process
- 5- a description of any prototype or working model;

In general, a manuscript that has been prepared for submission to a journal will satisfy this requirement.

5. Workable Extent/Scope [Describe the future course of related work, and possible variations of the present invention in terms of the broadest scope expected to be operable; if a **compound**, describe substitutions, breadth of substituents, derivatives, salts etc., if **DNA or other biological material**, describe modifications that are expected to be operable, if a **machine or device**, describe operational parameters of the device or a component thereof, including alternative structures for performing the various functions of the machine or device]

The future scope of this work could entail (1) Use of novel Hedgehog inhibitors to investigate their potential efficacy in inhibiting the growth of SCLC, and (2) Use of cyclopamine or other novel Hedgehog inhibitors to treat SCLC in humans.

6. Key Words [Please list specific key words that accurately describe the present invention]

Lung Cancer, Hedgehog pathway, Hedgehog pathway inhibitors, Treatment

7. References [Please list the closest and most relevant journal citations, patents, general knowledge or other public information related to the invention]

- Watkins DN et al, Hedgehog signaling within airway epithelial progenitors and in small cell lung cancer. Nature 2003 *in press*
- Berman DM et al, Medulloblastoma Growth Inhibition by Hedgehog Pathway Blockade. Science, 2002; 297:1559-1561
- Taipale J et al, Effects of oncogenic mutations in Smoothened and Patched can be reversed by cyclopamine. Nature 2000; 6799:1005-1009
- Cooper et al, Teratogen-Mediated Inhibition of Target Tissue Response to Shh Signaling. Science, 1998; 280:1603-1607.

SOFTWARE IMPLEMENTATION OF THE INVENTION

Indicate if this disclosure of invention is software or if software is implemented in the invention.

☐ Yes ☒ No

If Yes, describe the implementation of the software completely, using the outline given below.

1. **Scope of Work** [Is the work original? Is it created within the scope of your employment at JHU? Please explain the circumstances of program's development]

2. **Software Developers** [Please list any developers of the software if different from invention]

None ☐

3. **Software Derivation** [If software is a derivative of an existing work, please explain the original work's source and the modification]

None ☐

4. **Third Party Content** [Identify any third party content or other elements and their source included in the software]

None ☐

5. **Brief Software Description** [Please characterize how robust and user friendly the work is.]

RESEARCH SUPPORT INFORMATION

Indicate ALL contributions to the development of the invention in terms of personnel, money, materials and facilities etc.

Check each funding source that applies to this invention:

☐ None ☒ Federal Sponsor(s) ☐ University Funding ☐ Commercial Funding ☒ Other

For each funding source, provide the below information. Additionally, if "Commercial" or "Other" Funding was used, please attach a copy of each such award notice.

<u>Granting/Funding Source</u>	<u>Award/Contract Number</u>	<u>Title of Grant</u>	<u>Copy Attached</u>
NCI SPORE	CA058184	Inhibition of Hedgehog Signaling in Human Lung Cancer	X
FAMRI	Cyclopamine Inhibits Hedgehog Signaling and Growth in Lung Cancer Cells		X
			<input type="checkbox"/>
			<input type="checkbox"/>
			<input type="checkbox"/>
			<input type="checkbox"/>

AGREEMENT SUPPORT INFORMATION

Were any materials, equipment or software under a Special Agreement, such as Material Transfer agreements, purchase agreements, sponsored research agreements, or the like used? ☐ Yes ☒ No

If yes, please provide the following information for each item and attach a copy of the Agreement.

<u>Source of Materials</u>	<u>Materials</u>	<u>Copy Attached</u>
		<input type="checkbox"/>
		<input type="checkbox"/>
		<input type="checkbox"/>
		<input type="checkbox"/>





ACKNOWLEDGMENT, CERTIFICATION and ASSIGNMENT OF INVENTION

In order for this Report of Invention to be complete and processed by LTD, it must be signed and dated by:

- (1) the JHU Department Director for each JHU department involved with the development of this invention (SECTION A), and,
- (2) ALL Inventors (SECTIONS B and C), including those Inventors not subject to The Johns Hopkins University Intellectual Property Policy. Each Inventor must complete only one of Sections B or C (See explanations below).
- (3) Please duplicate Sections A, B and/or C as needed for proper completion with ALL appropriate signatures.

SECTION A. JHU DEPARTMENT DIRECTOR'S ACKNOWLEDGEMENT

I have read and understood this Report of Invention.

 JHU Department Director Signature	 Typed or Printed Name  JHU School / Department	 Date
JHU Department Director Signature	Typed or Printed Name JHU School / Department	Date
JHU Department Director Signature	Typed or Printed Name JHU School / Department	Date





SECTION B. JHU INVENTOR CERTIFICATION and ASSIGNMENT

This section is to be completed only by those JHU personnel subject to The Johns Hopkins University Intellectual Property Policy. Non-JHU Inventors, and HHMI or KKI Inventors at JHU are subject to a separate assignment and must complete Section C. JHU Inventors who believe they are not subject to The Johns Hopkins University Intellectual Property Policy for the invention described herein must complete Section C.

I/we, the Inventors, hereby certify that the information set forth in this Report of Invention is true and complete to the best of my/our knowledge.

I/we, the Inventors, hereby certify that I/we will promptly advise LTD of any commercial interest regarding the invention described herein.

I/we, the Inventor(s), subject to The Johns Hopkins University Intellectual Property Policy and not under an obligation to assign intellectual property rights to another party, hereby affirm that in consideration for The Johns Hopkins University's evaluation of commercial potential and a share of income which I/we may receive upon commercialization of my/our invention, on the date of my/our signature(s) as indicated below do hereby assign and transfer my/our entire right, title and interest in and to the invention described herein unto The Johns Hopkins University, its successors, legal representatives and assigns.

	D Neil Watkins MBBS PhD	3/7/03
JHU Inventor Signature	Typed or Printed Name	Date
	David M Berman MD PhD	3/7/03
JHU Inventor Signature	Typed or Printed Name	Date
	Philip A Beachy PhD	3/7/03 05-28-03. 9P Invalid signature removed
JHU Inventor Signature	Typed or Printed Name	Date
	Stephen B Baylin MD	3/7/03
JHU Inventor Signature	Typed or Printed Name	Date
JHU Inventor Signature	Typed or Printed Name	Date


SECTION C. INVENTOR CERTIFICATION without Assignment

Non-JHU Inventors, and HHMI or KKI Inventors at JHU must complete this Section.

Any JHU Inventor(s) who believes themselves not subject to assignment via The Johns Hopkins University Intellectual Property Policy for the specific invention described herein (e.g. the invention was made under a consulting agreement or other) must sign below and summarize the reasons for their belief. LTD will review this summary and may, depending upon the unique facts of the case, require assignment of the invention at a future date.

I/we, the Inventor(s), hereby certify that the information set forth in this Report of Invention is true and complete to the best of my/our knowledge.

I/we, the Inventor(s) who are HHMI or KKI inventors at JHU, hereby certify that I/we will promptly advise LTD of any commercial interest regarding the invention described herein.

 Inventor Signature	Philip Beachy Typed or Printed Name	3/7/03 Date
Exception to assignment via the JHU Intellectual Property Policy: <input checked="" type="checkbox"/> HHMI, <input type="checkbox"/> KKI, <input type="checkbox"/> Non-JHU personnel, <input type="checkbox"/> Other If Other, please provide details of the exception:		
 Inventor Signature	 Typed or Printed Name	 Date
Exception to assignment via the JHU Intellectual Property Policy: <input type="checkbox"/> HHMI, <input type="checkbox"/> KKI, <input type="checkbox"/> Non-JHU personnel, <input type="checkbox"/> Other If Other, please provide details of the exception:		
 Inventor Signature	 Typed or Printed Name	 Date
Exception to assignment via the JHU Intellectual Property Policy: <input type="checkbox"/> HHMI, <input type="checkbox"/> KKI, <input type="checkbox"/> Non-JHU personnel, <input type="checkbox"/> Other If Other, please provide details of the exception:		

Please submit reports to:

Licensing and Technology Development
Johns Hopkins University
111 Market Place, Suite 906
Baltimore, Maryland 21202
Phone: (410) 347-3222 Fax: (410) 347-3201

Hedgehog signalling within airway epithelial progenitors and in small-cell lung cancer

D. Neil Watkins*, David M. Berman†‡, Scott G. Burkholder*, Baolin Wang‡, Phillip A. Beachy‡ & Stephen B. Baylín*

* Sidney Kimmel Comprehensive Cancer Center, † Department of Pathology, ‡ Department of Molecular Biology and Genetics, and Howard Hughes Medical Institute, Johns Hopkins University School of Medicine, Baltimore, Maryland 21231, USA

Embryonic signalling pathways regulate progenitor cell fates in mammalian epithelial development and cancer^{1,2}. Prompted by the requirement for sonic hedgehog (Shh) signalling in lung development^{3,4}, we investigated a role for this pathway in regeneration and carcinogenesis of airway epithelium. Here we demonstrate extensive activation of the hedgehog (Hh) pathway within the airway epithelium during repair of acute airway injury. This mode of Hh signalling is characterized by the elaboration and reception of the Shh signal within the epithelial compartment, and immediately precedes neuroendocrine differentiation. We reveal a similar pattern of Hh signalling in airway development during normal differentiation of pulmonary neuroendocrine precursor cells, and in a subset of small-cell lung

cancer (SCLC), a highly aggressive and frequently lethal human tumour with primitive neuroendocrine features. These tumours maintain their malignant phenotype *in vitro* and *in vivo* through ligand-dependent Hh pathway activation. We propose that some types of SCLC might recapitulate a critical, Hh-regulated event in airway epithelial differentiation. This requirement for Hh pathway activation identifies a common lethal malignancy that may respond to pharmacological blockade of the Hh signalling pathway.

Sonic hedgehog (Shh), a mammalian hedgehog (Hh) pathway ligand, mediates epithelial–mesenchymal interactions in lung development by signalling to adjacent lung mesenchyme, as indicated by expression of the Hh receptor and pathway target *Patched* (*Ptch*)⁵. Loss of Shh function results in severe lung defects associated with failure of branching morphogenesis^{3,4}. As developmental pathways regulate progenitor cell fates and differentiation in some regenerating mammalian epithelia^{1,2}, we hypothesized that Hh signalling might be important in airway epithelial repair.

In contrast to the skin and colon, adult airway epithelium rarely proliferates unless injured⁶. To uncover a role for Hh signalling in this process, we studied a mouse model of acute airway repair in which Clara cells, specialized airway epithelial cells predominant in distal conducting airways, are depleted within 24 h of systemic naphthalene administration⁶. Activation of a putative airway progenitor results in epithelial regeneration within three days, with increased numbers of airway neuroendocrine cells—a normally rare cell type implicated in the regulation of airway epithelial growth and




Figure 1 Hedgehog signalling in airway repair and development. **a**, Immunohistochemical detection of Shh and Gli1 in adult mouse airways is negative in normal airways (left panels), but positive for both Shh and Gli1 in serial sections 3 days after naphthalene injury (middle panels). By 4 days after naphthalene treatment (right panels), Gli1-positive cells are reduced in number (arrow). Serial sections demonstrate that nascent

CGRP-positive cells do not express stained Gli1. Scale bar, 50 μ m. **b**, Quantitative analysis of bronchial epithelial staining in **a** ($n = 4$, mean \pm s.e.m.). **c**, Shh signalling in E13.5 lungs. Shh immunostaining in embryonic airway epithelium is shown in the left panel. The right panel shows X-gal staining of lungs obtained from E13.5 *Ptch-LacZ* mouse embryos, demonstrating intense mesenchymal staining. Scale bar, 25 μ m. **d**, Clusters of LacZ-positive cells (arrows) are seen in the airway epithelium of E16.5 (left panel) and adult (right panel) mice. Scale bar, 25 μ m. **a**, airway; **m**, mesenchyme; **bm**, basement membrane. **e**, Confocal immunofluorescence detection of Hh signalling in lung development. The top row demonstrates expression of both CGRP and *Ptch* in an E16.5 airway (arrow), similar to that shown in **d**. The bottom row shows expression of CGRP (arrow) adjacent to Shh-expressing epithelial cells (see high-magnification inset). Scale bar, 25 μ m.

development^{6,7}. In regenerating airways, we observed marked expression of both Shh ligand and Gli1, a transcriptional target of Hh signalling⁸, in the epithelial compartment 72 h after naphthalene injury (Fig. 1a). By day 4, Gli1 was not observed in nascent airway epithelial cells expressing calcitonin gene-related peptide (CGRP), a marker of neuroendocrine differentiation (Fig. 1a, b). These data show that acute airway epithelial regeneration results in widespread activation of airway intraepithelial Hh signalling, which immediately precedes neuroendocrine differentiation.

Embryonic lung epithelial cells express Shh, which is thought to signal to adjacent lung mesenchyme to regulate branching morphogenesis³⁻⁵. In light of this, our detection of Shh and Gli1 within the epithelial compartment during airway epithelial regeneration was unexpected. To determine whether such intraepithelial signalling occurred in embryonic lung development, we studied mice in which one copy of *Ptch* is replaced in-frame with the β -galactosidase (β -gal) gene by homologous recombination⁹. As *Ptch* is a transcriptional target of the Gli proteins, expression of β -gal indicates activation of the Hh pathway^{9,10}. Early gestation (embryonic day (E)13.5) embryos showed expression of Shh protein in the primitive lung endoderm, and intense β -gal expression in the adjacent mesenchyme (Fig. 1c). By contrast, later lung development (E16.5) was characterized by clusters of β -gal-expressing cells in the developing airway epithelium (Fig. 1d). Small numbers of cells expressing β -gal persist in the basal layer of the adult bronchial epithelium (Fig. 1d). Similar clusters of epithelial cells expressing the neuroendocrine marker CGRP and *Ptch* were observed by confocal immunofluorescence in E16.5 airways, immediately adjacent to cells expressing Shh (Fig. 1e). These data suggest that during normal development, neuroendocrine precursors within the airway epithelial compartment respond to a Shh signal elaborated by adjacent airway epithelial cells.

SCLC is an aggressive, highly lethal malignancy with primitive neuroendocrine features¹¹. As aberrant reactivation of developmental pathways may have a role in cancer growth^{1,2}, we wondered whether the epithelial Hh signalling we had observed in airway embryogenesis and repair might persist in SCLC. Analysis of SCLC tissue showed that five out of ten tumours expressed both Shh and Gli1 (Fig. 2a; see also Supplementary panel a). Out of 40 non-SCLC (NSCLC) tumours, nine demonstrated Shh expression and of these, four demonstrated co-expression of Gli1 (Fig. 2a; see also Supplementary panel a). These data provide indirect evidence of persistent activation of Hh signalling in lung cancer, predominantly in SCLC. These findings were confirmed by analysis of human lung cancer cell lines. Notably, all seven SCLC and seven NSCLC cell lines expressed Shh protein (Fig. 2b). Out of five breast and eight colon cancer cell lines examined, only one (CACO2) expressed Shh protein, and none expressed Gli1 protein, as shown by western blot analysis (data not shown). Importantly, expression of both Shh and Gli1 proteins was observed in five out of seven SCLC lines, and this correlated with increased expression of *Ptch* messenger RNA (Fig. 2b). In contrast, NSCLC lines expressed Shh and low levels of *Ptch*, but not Gli1. These data are summarized in Supplementary panel b.

To determine how Hh signalling might function in these tumours, we co-cultured cancer cells with Shh-LIGHT2 cells, a fibroblast reporter cell line that responds to exogenous Shh by activation of an integrated Gli-responsive luciferase reporter¹⁰. Some NSCLC cells that express Shh are capable of heterologous cell signalling to the reporter cell line (Fig. 2c), suggesting that NSCLC retains the Shh export properties of primitive lung endodermal cells that signal to adjacent mesenchymal cells in early development. By contrast, the SCLC cells we studied display a marked reduction in this ability to signal to adjacent cells. These




Figure 2 Hh signalling in lung cancer. **a**, Examples of Shh and Gli1 immunostaining in human lung cancer tissue. Note the widespread co-expression of Shh and Gli1 in SCLC, which is reduced in the NSCLC example. **b**, Expression of Hh signalling components in lung cancer cell lines. The top panel shows immunoblotting (IB) data for expression of Shh, Gli1 and GAPDH. The bottom panel demonstrates *Ptch* mRNA expression in the same cell lines detected by RNase protection assay (RPA). The markers along the right indicate relative molecular mass. **c**, Induction of *Gli*-luciferase activity in Shh-LIGHT2 reporter cells

co-cultured with purified Shh-Np or the cell lines indicated on the x axis. Luciferase activity is normalized to a Renilla luciferase internal control ($n = 6$, mean \pm s.e.m.). **d**, Shh and Gli1 expression in NCI-H249 SCLC xenograft cells detected by dual-label immunohistochemistry. Brown, Shh; red, Gli1. The left panel shows a tumour cell expressing Shh alone (arrow); the right panel shows a Shh-expressing tumour cell (top arrow) and an adjacent Gli1-expressing tumour cell (bottom arrow).

data demonstrate that distinct types of lung cancer cells recapitulate different aspects of Shh signalling seen in lung development and repair.

We next addressed the mechanism of Hh pathway activation in SCLC. Dual-label immunostaining for Shh and Gli1 in SCLC nude mouse xenografts demonstrated Shh-expressing cells adjacent to Gli1-expressing cells (Fig. 2d). These data suggest juxtacrine Hh pathway activation in SCLC markedly similar to that observed in airway development and repair. Next, we asked whether ligand-driven Hh pathway activation promotes growth of SCLC. Inhibition of Shh ligand activity in NCI-H249 and NCI-H1618 SCLC cells with the 5E1 Shh-N monoclonal antibody¹² resulted in growth inhibition (Fig. 3a). Although NCI-H157 NSCLC cells express Shh, they do not express Gli1 protein, and are not affected by 5E1 treatment (Fig. 3a). These data demonstrate that growth of SCLC cells *in vitro* is dependent on ligand-mediated activation of the Hh pathway, and suggest the presence of a normal Ptch receptor, confirmed by sequencing of *Ptch* complementary DNA in both NCI-H249 and NCI-H1618 SCLC cells generated by reverse transcription-polymerase chain reaction (RT-PCR) (data not shown).

The *Veratrum* alkaloid cyclopamine specifically inhibits the Hh pathway^{10,13,14} through interaction with the Hh signalling protein smoothened^{15,16}. Moreover, cyclopamine blocks the oncogenic effects of mutations of *Ptch* in fibroblasts¹⁰, and inhibits the malignant growth of medulloblastoma cells lacking *Ptch* function¹⁷. Treatment of NCI-H249 SCLC cells with cyclopamine, or a more potent analogue KAAD-cyclopamine¹⁰, resulted in significant growth inhibition, whereas tomatidine, a closely related compound that lacks the capacity to inhibit Hh signalling, had no effect (Fig. 3b). The effects of cyclopamine and KAAD-cyclopamine on the growth of SCLC reflect their relative potency in silencing Hh pathway activation *in vitro*¹⁰. None of KAAD-cyclopamine, cyclopamine or tomatidine was able to affect growth of NCI-H157 NSCLC cells (Fig. 3c). The growth-inhibitory effect of cyclopamine, if due to Hh pathway blockade, should be bypassed by constitutive overexpression of the Hedgehog pathway effector Gli1 (ref. 17). We indeed observed that stable expression of a Flag-tagged Gli1 protein¹⁸ protected NCI-H249 SCLC cells from growth inhibition by cyclopamine, whereas a Gli1 mutant lacking the zinc finger DNA-binding domain had no effect (Fig. 3d). Treatment of nine cancer cell lines with cyclopamine at concentrations up to 10 μ M demonstrated growth inhibition only in SCLC cells that expressed both Shh and its transcriptional effector Gli1 (Supplementary panel b). These data show that cyclopamine induces growth inhibition in SCLC cells expressing both Shh and Gli1 by specific inhibition of the Hh pathway.

We next investigated the relationship between Hh pathway blockade by cyclopamine and growth arrest in SCLC. Unsynchronized NCI-H249 SCLC cells treated with 5 μ M cyclopamine for 72 h demonstrated arrest of the cell cycle in G0/G1 (Fig. 3e) and apoptosis indicated by an increase in cleaved PARP (Fig. 3f). Analysis of *Ptch* mRNA expression revealed downregulation in response to cyclopamine treatment (Fig. 3g). These results indicate silencing of Hh pathway activation at concentrations of cyclopamine that induce both growth arrest and apoptosis. We next investigated the possibility that SCLC cells might express transcripts indicative of a progenitor cell phenotype. We detected expression of BMP4, a morphogen and putative target of Hh expressed in lung epithelial embryogenesis¹⁹, and nestin, an intermediate filament characteristic of neural stem cells in medulloblastoma¹⁷ (Fig. 3h). Treatment of NCI-H249 SCLC cells with cyclopamine for 48 h inhibited expression of both these genes (Fig. 3h), as well as the expression of human ASH-1, a transcription factor required for pulmonary neuroendocrine differentiation²⁰. These changes in gene expression suggest that Hh signalling maintains a progenitor cell fate in SCLC.

Pathological activation of Hh signalling is associated with

medulloblastoma, a malignant brain tumour thought to arise from the granule cells of the cerebellum^{9,21}. Maintenance of abnormal progenitor-like fates through continued Hh pathway activation is essential for malignant growth of these tumours *in vivo*¹⁷. We




Figure 3 Hh pathway activation is essential for the growth of SCLC. **a**, Growth of cancer cell lines treated with monoclonal antibodies against β -galactosidase (β -gal) as a control, or Shh for 4 days. **b**, NCI-H249 SCLC cell growth after 5 days, treated with tomatidine, cyclopamine or KAAD cyclopamine at the indicated concentrations. **c**, Identical experiment performed in NCI-H157 NSCLC cells. **d**, Response of stably transfected NCI-H249 SCLC cells to treatment with cyclopamine when expressing neomycin resistance (Neo), a mutant Gli1 lacking the zinc finger domain (Flag-Gli1ZFD), Gli1 (Flag-Gli1), and wild-type untransfected (WT) cells. Cell viability was measured by MTT assay, detected at an absorbance at 540 nm (A_{540}) ($n = 6$) and expressed as a percentage of control \pm s.e.m. **e**, Cell cycle analysis in NCI-H249 cells treated with tomatidine or cyclopamine (5 μ M). Percentages in each phase of the cell cycle are shown below and are shown as the mean of three experiments. **f**, Cleaved PARP expression in NCI-H249 and NCI-H417 SCLC cells treated with tomatidine (-) or cyclopamine (+) (5 μ M). **g**, *Ptch* mRNA expression in NCI-H249 SCLC cells detected by northern blot analysis after treatment with cyclopamine. 28s RNA stained with ethidium bromide is shown as a loading control. **h**, RT-PCR analysis of transcripts in NCI-H249 SCLC cells. Cont, control; Tom, tomatidine treated; Cyc, cyclopamine treated.




Figure 4 Cyclopamine inhibits SCLC tumorigenicity. **a**, The top panel shows soft agar growth of NCI-H249 SCLC cells treated with cyclopamine. Plates were stained with ethidium bromide. The bottom panel shows colony formation of NCI-H249 SCLC cells treated with cyclopamine (5 μ M) and stably transfected with neomycin resistance (Neo^r), mutant Gli1 (Flag-Gli1ZFD) or Gli1 (Flag-Gli1). **b**, Quantitative analysis of the experiment described in **a**. Data are shown as mean colonies per well \pm s.e.m. ($n = 6$). **c**, Growth of NCI-H249 nude mouse subcutaneous xenografts in animals treated with tomatidine or cyclopamine for 10 days. **d**, Identical experiment to that shown in **c** except that A549 NSCLC cells were used. Data are shown as mean tumour volume \pm s.e.m. as a percentage of tumour volume at day 0 ($n = 7$).

wondered whether SCLC cells were similarly dependent on Hh signalling for their malignant behaviour. NCI-H249 SCLC cells treated with cyclopamine showed reduced soft agar clonogenicity—an *in vitro* assay of tumorigenicity (Fig. 4a, b). This effect was reversed in cells overexpressing the Hh pathway transcriptional effector Gli1 (Fig. 4a, b). We next tested the ability of systemic cyclopamine treatment to inhibit the growth of SCLC xenografts in nude mice. Mice bearing xenografts were treated subcutaneously with 25 mg kg⁻¹ day⁻¹ cyclopamine as described¹⁷. Growth inhibition was observed in three SCLC lines: NCI-H249 (Fig. 4c), as well as NCI-H417 and NCI-H1618 (data not shown). No effect was observed in A549 NSCLC cells (Fig. 4d) nor in HCT-116 colon cancer xenografts (data not shown). These data are summarized in Supplementary panel b, and show that Hh signalling is required for the growth *in vivo* of SCLC cells that express both Shh and Gli1.

We have shown that Hh signalling in airway epithelium is not limited to epithelial-mesenchymal interactions, but can be contained within the airway epithelial compartment during embryonic neuroendocrine differentiation and airway repair. Taking evidence that links Hh signalling to cerebellar progenitor cell differentiation into consideration^{21–23}, we propose a similar role for this pathway in the regulation of airway progenitor cell fates, which may be specified immediately before the divergence of neuroendocrine and non-neuroendocrine lineages. The dependency of SCLC cells on Hh pathway activation is also notable in that it relies on the presence of Shh ligand, it occurs in the absence of mutations in *Ptch*, and recapitulates juxtacrine Hh signalling seen in development and airway repair. SCLC may represent a malignancy arising from an airway epithelial progenitor that retains both Hh signalling and primitive features of pulmonary neuroendocrine differentiation. The vulnerability of SCLC to Hh pathway blockade may represent a new therapeutic approach to a disease with a poor prognosis²⁴. □

Methods

Detection of β -gal expression

Ptch-LacZ mice were maintained and genotyped as described⁹. 5-bromo-4-chloro-3-indolyl- β -D-galactoside (X-gal) staining in microdissected mouse lungs was performed overnight as described²⁵, followed by post-fixation in formalin, paraffin embedding and sectioning. We used wild-type littermates as negative controls.

Immunohistochemistry

Single-colour DAB-immunoperoxidase staining was performed using a modification of the DAKO CSA system. Detailed protocols are available on request. Antibodies were from Santa Cruz Biotechnologies: Shh (N-19; sc-1194); Gli1 (N-16; sc-6153); CGRP (N-20; sc-8856). Shh, Ptch and Gli1 staining was optimized on paraffin sections from Shh wild-type and knockout embryos. Gli1 staining was further confirmed in Flag-Gli1-overexpressing Cos-7 cells by immunofluorescence. Peptide competition ablated staining in tumour samples and embryos. Dual-colour immunohistochemistry was performed using the DAKO Envision system. Dual-colour immunofluorescence was performed on fresh-frozen sections fixed in paraformaldehyde using Molecular Probes Alexa secondary antibodies.

Western immunoblot

Whole-cell lysates were sonicated in 2% SDS/50 mM TrisHCl, pH 8. Western blot using rabbit polyclonal antibodies for Shh-N were performed as described²⁶. A rabbit polyclonal antibody to Gli1 was developed as described²⁷ using a glutathione S-transferase fusion protein containing amino acid residues 216–271 of human Gli1. Anti-cleaved PARP was obtained from Promega.

RNAse protection assay and northern blot analysis

RNAse protection assay (RPA) was performed as described²⁸ using a *Ptch*-specific antisense RNA probe corresponding to bases 1338–1788 of the human patched-1 cDNA (GLI1335863) generated by RT-PCR and subcloned into pCR-TOPOII (Stratagene). Northern blotting of 10 μ g total RNA was performed as described²⁹, and probed with a *Ptch*-specific cDNA probe obtained from the same construct.

RT-PCR

Total cellular RNA was treated with DNase, reverse transcribed, and amplified for 31 cycles at an annealing temperature of 55 °C. Primers used were BMP4(+), 5'-CTTTACCGGC TTGAGTCTGGG-3'; BMP4(-), 5'-CCCAATTCCCACTCCCTTGAG-3'; GAPDH(+), 5'-ATCTTCCAGGAGCCAGATCCC-3'; GAPDH(-), 5'-CGTTCGGCTCAGGGATGA CCT-3'; ASH-1(+), 5'-CCATGCAAGCTCTGCCAAG-3'; ASH-1(-), 5'-TGACC AACTTGACGCGTTGC-3'; nestin(+), 5'-CTCTGGAGAGGAGATCAAG-3'; nestin(-), 5'-CCTTTGTCAGAGCTCTCAGT-3'.

Shh-LIGHT2 reporter assay

Superconfluent reporter cells were cultured as described¹⁰, then co-cultured in low serum conditions in the presence of 1×10^5 cells per well of the cell line of interest or purified Shh-Np¹⁰. Luciferase and Renilla luciferase assays were performed using the Promega Dual Luciferase Reporter Assay system.

Cell culture experiments

Cell lines were obtained from American Type Culture Collection (ATCC). Shh inhibitor experiments were performed in 0.5% calf serum. Cyclopamine was obtained from Toronto Research Chemicals. Both were dissolved as $\times 1,000$ stocks in DMSO medium. Flag-tagged Gli1 vectors were obtained from the Joyner laboratory¹⁴. To generate NCI-H249 SCLC cells overexpressing each of the Gli vectors, mass cultures were stably co-transfected using lipofectamine (Invitrogen) with the Flag-Gli vector of interest, and pcDNA3.1 (Stratagene) to confer neomycin resistance. 5E1 anti-ShhN monoclonal antibody was used at a concentration of $10 \mu\text{g ml}^{-1}$ as described¹⁵. Soft agar assays were performed as described¹⁰. Cells were seeded into six-well plates at a density of 20,000 cells per well in agar containing 2% calf serum. MTT assays were performed as described¹⁰.

Nude mouse xenografts

Tumour cell lines were injected subcutaneously at 1×10^7 cells per mouse and allowed to grow to a maximum diameter of 5 mm. Cyclopamine was administered as described¹⁷. Tumours were measured daily and the tumour volume calculated as described¹⁰.

Received 1 November 2002; accepted 12 February 2003; doi:10.1038/nature01493.

Published online 5 March 2003.

1. Reyes, T., Morrison, S. J., Clarke, M. F. & Weissman, I. L. Stem cells, cancer, and cancer stem cells. *Nature* 414, 105–111 (2001).
2. Taipale, J. & Beachy, P. A. The Hedgehog and Wnt signalling pathways in cancer. *Nature* 411, 349–354 (2001).
3. Piccoli, C. V., Lewis, P. M. & McMahon, A. P. Sonic hedgehog regulates branching morphogenesis in the mammalian lung. *Curr. Biol.* 8, 1083–1086 (1998).
4. Litingtung, Y., Lei, L., Westphal, H. & Chiang, C. Sonic hedgehog is essential for foregut development. *Nature Genet.* 20, 58–61 (1998).
5. Bellusci, S. et al. Involvement of Sonic hedgehog (Shh) in mouse embryonic lung growth and morphogenesis. *Development* 124, 53–63 (1997).
6. Reynolds, S. D., Giangreco, A., Power, J. H. & Stripp, B. R. Neuroepithelial bodies of pulmonary airways serve as a reservoir of progenitor cells capable of epithelial regeneration. *Am. J. Pathol.* 156, 269–278 (2000).
7. Peake, J. L., Reynolds, S. D., Stripp, B. R., Stephens, K. E. & Pinkerton, K. E. Alteration of pulmonary neuroendocrine cells during epithelial repair of naphthalene-induced airway injury. *Am. J. Pathol.* 156, 279–286 (2000).
8. Lee, J., Platt, K. A., Censullo, P. & Ruiz i Altaba, A. Gli1 is a target of Sonic hedgehog that induces ventral neural tube development. *Development* 124, 2537–2552 (1997).
9. Goodrich, L. V., Milenkovic, L., Higgins, K. M. & Scott, M. P. Altered neural cell fates and medulloblastoma in mouse patched mutants. *Science* 277, 1109–1113 (1997).
10. Taipale, J. et al. Effects of oncogenic mutations in Smoothened and Patched can be reversed by cyclopamine. *Nature* 406, 1005–1009 (2000).
11. Zochbauer-Müller, S., Gazdar, A. F. & Minna, J. D. Molecular pathogenesis of lung cancer. *Annu. Rev. Physiol.* 64, 681–708 (2002).
12. Ericson, J., Morton, S., Kawakami, A., Roelink, H. & Jessell, T. M. Two critical periods of Sonic Hedgehog signaling required for the specification of motor neuron identity. *Cell* 87, 661–673 (1996).
13. Incardona, J. P., Gaffield, W., Kapur, R. P. & Roelink, H. The teratogenic Veratrum alkaloid cyclopamine inhibits sonic hedgehog signal transduction. *Development* 125, 3553–3562 (1998).
14. Cooper, M. K., Porter, J. A., Young, K. E. & Beachy, P. A. Teratogen-mediated inhibition of target tissue response to Shh signaling. *Science* 280, 1603–1607 (1998).
15. Chen, J. K., Taipale, J., Cooper, M. K. & Beachy, P. A. Inhibition of Hedgehog signaling by direct binding of cyclopamine to Smoothened. *Genes Dev.* 16, 2743–2748 (2002).
16. Chen, J. K., Taipale, J., Young, K. E., Maiti, T. & Beachy, P. A. Small molecule modulation of Smoothened activity. *Proc. Natl Acad. Sci. USA* 99, 14071–14076 (2002).
17. Berman, D. M. et al. Medulloblastoma growth inhibition by Hedgehog pathway blockade. *Science* 297, 1559–1561 (2002).
18. Park, H. L. et al. Mouse Gli1 mutants are viable but have defects in SHH signaling in combination with a Gli2 mutation. *Development* 127, 1593–1605 (2000).
19. Weaver, M., Yingling, J. M., Dunn, N. R., Bellusci, S. & Hogan, B. L. Bmp signaling regulates proximal-distal differentiation of endoderm in mouse lung development. *Development* 126, 4005–4015 (1999).
20. Borges, M. et al. An achaete-scute homologue essential for neuroendocrine differentiation in the lung. *Nature* 386, 852–855 (1997).
21. Kenney, A. M. & Rowitch, D. H. Sonic hedgehog promotes G(1) cyclin expression and sustained cell cycle progression in mammalian neuronal precursors. *Mol. Cell Biol.* 20, 9055–9067 (2000).
22. Dahmane, N. & Ruiz-i-Altaba, A. Sonic hedgehog regulates the growth and patterning of the cerebellum. *Development* 126, 3089–3100 (1999).
23. Wechsler-Reya, R. J. & Scott, M. P. Control of neuronal precursor proliferation in the cerebellum by Sonic Hedgehog. *Neuron* 22, 103–114 (1999).
24. Johnson, D. H. Management of small cell lung cancer: current state of the art. *Chest* 116, S25S–S30S (1999).
25. Hogan, B. L., Beddington, R., Constantini, F. & Lacy, E. *Manipulating the Mouse Embryo* (Cold Spring Harbor Press, Plainview, 1994).
26. Chang, D. T. et al. Products, genetic linkage and limb patterning activity of a murine hedgehog gene. *Development* 120, 3339–3353 (1994).
27. Wang, B., Fallon, J. F. & Beachy, P. A. Hedgehog-regulated processing of Gli3 produces an anterior/posterior repressor gradient in the developing vertebrate limb. *Cell* 100, 423–434 (2000).
28. Sriuranpong, V. et al. Notch signaling induces rapid degradation of achaete-scute homolog 1. *Mol. Cell Biol.* 22, 3129–3139 (2002).
29. Nakakura, E. K. et al. Mammalian Scratch: a neural-specific Snail family transcriptional repressor. *Proc. Natl Acad. Sci. USA* 98, 4010–4015 (2001).
30. Park, B. H., Vogelstein, B. & Kinzler, K. W. Genetic disruption of PPARdelta decreases the tumorigenicity of human colon cancer cells. *Proc. Natl Acad. Sci. USA* 98, 2598–2603 (2001).

Supplementary information accompanies the paper on Nature's website (<http://www.nature.com/nature>).

Acknowledgements We acknowledge M. Scott for the donation of the *Ptch-LacZ* mice; A. Joyner for the Flag-Gli1 expression vectors; J. Chen for the gift of KAAD cyclopamine; J. Taipale, S. Karhadkar, B. Nelkin and K. Schuebel for discussions; and E. Gabrielson for help in obtaining lung cancer tissue. We also thank K. Young and L. Metzler and the Sidney Kimmel Comprehensive Cancer Center Cell Imaging Facility for technical assistance. This work is supported by the Flight Attendant Medical Research Institute and the NCI/SPORC. P.A.B. is an investigator of the Howard Hughes Medical Institute.

Competing Interests statement The authors declare competing financial interests: details accompany the paper on Nature's website (<http://www.nature.com/nature>).

Correspondence and requests for materials should be addressed to D.N.W. (e-mail: mwatkins@jhmi.edu).

All publications, patents and patent applications disclosed herein are incorporated into this application by reference in their entirety.

For example: "Sambrook *et al.*, Molecular Cloning, A Laboratory Manual (volumes I-III) 1989, Cold Spring Harbor Laboratory Press, USA", "Harlowe and Lane, Antibodies a Laboratory Manual 1988 and 1998, Cold Spring Harbor Laboratory Press, USA" and "Ausubel *et al.*, Current Protocols 2001, John Wiley and sons, Inc." provide sections describing methodology for antibody generation and purification, diagnostic platforms, cloning procedures, etc. that may be used in the practice of the instant invention.

17